

## Optimized Lifestyle via Timing of Food Intake: Bridging Ruminant Agriculture to Human Health

Akbar Nikkhah\*<sup>1</sup><sup>1</sup>Chief Highly Distinguished Professor, Department of Animal Sciences, Faculty of Agricultural Sciences, University of Zanja, Iran

\*Corresponding author: Dr. Akbar Nikkhah, Chief Highly Distinguished Professor, Foremost Principal Highly Distinguished Elite-Generating Scientist, Department of Animal Sciences, Faculty of Agricultural Sciences, University of Zanja, National Elite Foundation, Iran, E-mail: anikkha@yahoo.com

Received: 06-02-2015

Accepted: 06-26-2015

Published: 07-25-2015

Copyright: © 2015 Akbar Nikkhah

### Abstract

Ruminant discoveries help construct a promise to develop family programs on timing of eating to improve cell metabolism and overall body health. In light of the increasing concerns on obesity, diabetes, cardiovascular complexities, and cancer, public education must adopt optimum times of nutrient intake over the 24-h period. Such a guideline would include avoiding large evening and night meals, consuming early post-exercise breakfast, and distributing frequent meals during morning and early afternoon hours. Timing of food intake is a feasible approach to improve life quality in the postmodern era.

**Key words:** Life quality; Timing of intake; Ruminant; Human health

### Introduction

This policy article aims to increase public awareness of the untraveled highway of 'eating timing' to optimizing integrative metabolism and health of the man in the new post-modern era. The most recent discoveries in comparative livestock, rat and human nutritional physiology have contributed to the emergence of the novel science of 'chronophysiology management'. This science is concerned with circadian rhythms of cell metabolism and how they should be synchronized with the external factors including but not limited to timing of eating and photoperiod [1-4].

### Discoveries, Intuitions and Discussion

Recent findings have revealed that ruminants as irreplaceable working models with sophisticated metabolism and physiology respond dramatically to timing of eating [5,6]. Dairy cows fed in evening vs. morning under thermoneutral

and controlled environments produced greater milk energy and in some cases consumed greater feed that is a sign of good metabolic health [6,7]. These findings were substantiated in following studies [8]. In addition, we do know that glucose tolerance and insulin sensitivity in human are impaired as day leaves for night [9,10]. The fact that evening fed cows produce more milk and may eat more feed stems from their considerably higher metabolic turnover of peripheral tissues, when compared to human. However, the unique brain metabolism in human must be contemplated differently when aiming to interpret animal data for interventions in human metabolism and medicine.

All in all, the science generated mainly over the last few decades established that when to eat matters more than much for public health. Insulin is not required very much in high-producing cows since they need not to deposit but instead to secrete nutrients out of the body. In other words, insulin action in high-producing ruminants is rather restrict-

ed to nutrient oxidation in fuelling splanchnic tissue's energy expenditure. In contrast, human metabolism is greatly dependent on and responsive to insulin all the time. Thus, in view of the evolutionary human metabolism, less glucose and thus less insulin have been required overnight when the body must rest and not highly metabolize. As a result, human has not evolved to develop a high-capacity insulin-dependent glucose and amino acid metabolism during evening and night times. As such, should food be greatly supplied overnight, when the body is not prepared to manage, impaired intermediary and integrative metabolism can expectedly occur.

### Implications

The integrative public education must concentrate on signifying the importance of timing of eating to ensure that internal physiology is properly managed to be adequately synchronized with the external environment. This can assure that people know and can feasibly practice how prevention is superior to treatment in overcoming metabolic disorders and various types of cancer. It is time to make science workable for all.

### Acknowledgments

The Ministry of Science Research and Technology, National Elite Foundation, and University of Zanjan, Iran, thanked for supporting the author's global programs of optimizing science edification in the third millennium.

### References

1. Nikkhah A. Eating time modulations of physiology and health: life lessons from human and ruminant models. *Iranian J Basic Med Sci.* 2012, 15(4): 787-794.
2. Nikkhah A. Ruminant chronophysiological management: an emerging bioscience. *Open Access Anim Physiol.* 2011, 3: 9-12.
3. Nikkhah A. Chronophysiology of ruminant feeding behavior and metabolism: an evolutionary review. *Biol. Rhythm Res.* 2013, 44(2): 197-218.
4. Nikkhah A. Bioscience of ruminant intake evolution: feeding time models. *Adv. Biosci. Biotechnol.* 2: 271-274.
5. Nikkhah A. Timing of feeding: a postmodern management strategy to modulate chronophysiological rhythms in rumen fermentation kinetics. *Biol Rhythm Res.* In press. 2014.
6. Nikkhah A, Furedi CJ, Kennedy AD, Crow GH, Plaizier JC. Effects of feed delivery time on feed intake, rumen fermentation, blood metabolites and productivity of lactating cows. *J Dairy Sci.* 2008, 91: 1-12.
7. Nikkhah A. *Time of Feeding an Evolutionary Science*, Lap Lambert Publishing, GmbH & Co. KG, Germany, 1978, 3260-32622.
8. Nikkhah A, C Furedi, A Kennedy, K Wittenberg, JC Plaizier. Feed delivery at 2100 h vs. 0900 h for lactating dairy cows. *Can. J. Anim. Sci.* 91(1): 113-122.
9. Nikkhah A. Eating timing and diabetes. *Int.J. Diabetol. Vas. Dis. Res.* 2014, 2:101.
10. la Fleur SE, Kalsbeek A, Wortel J, Fekkes MI, Buijs RM. A daily rhythm in glucose tolerance: a role for the suprachiasmatic nucleus. *Diabetes.* 50: 1237-1243.